

ORAL Abstract

How lifeguards would know the severity, treatment and outcome of drowning on the accident site?

Principal Author: Dr David Szpilman, MD

Medical Staff of Drowning Resuscitation Center and Aeromedical Helicopter - Fire Department of Rio de Janeiro (CBMERJ) – GMAR; Head of Adult Intensive Care Unit - Hospital Municipal Miguel Couto; Member of Brazilian National Resuscitation Council; Founder, Former President and Medical Director of Brazilian Life Saving Society – SOBRASA; BOD Member and Medical Committee of International Life-Saving Federation; BOD Member of Sports Medical Council;

Abstract

In a well established beach lifesaving service, respiratory or cardiopulmonary arrests occur in only 0.5% of all rescues and in 10% from those who need medical attention (1). Although there are some controversial issues, it is well established on these cases the need for respiratory support alone or cardiopulmonary resuscitation, assistance of a medical doctor at the accident site, and hospitalization. But one of the most difficult and challenging lifeguard decisions for those who provide Basic Life Support – BLS or Advanced Life Support – ALS at the rescue site is how to recognize the severity of the drowning victims with less life-threatening conditions appropriately and consequently be confident in front of the critical public to do the right intervention. Our purpose is to reassure lifeguards confidence in front of the public, establishing simple recognition signs to let them know the drowning severity, treatment and outcome.

Methods: We reviewed literature (Medline, references from previous articles, review articles and books and ILS Policies) searching for the words drowning and severity. All data which does not focus on rescue site were excluded. We focused our search to answer 3 questions: How to recognize drowning severity? How to treat according to severity? And when a lifeguard should seek medical advice or support?

Results: Two large researches were reviewed with a total of 87,339 rescues (1,2). From those, 3,234(3.7%) cases needed medical attention at the rescue site, but severity evaluation was based on 2,678 cases. Medical attendance was available in 100% of cases and was responsible for treatment and for triage to release home or for hospitalization. Drowning severity, treatment decision, outcome/mortality and the need to seek for medical advice or support is summarized in table 1.

Discussion: One of the most difficult and challenging tasks a lifeguard must face is the giving of first aid. This part of his or her responsibilities requires lengthy theory and practical training exercises using many different situations. This is always done by the lifeguard, alone or with a partner, in front of the public who often interrupt with conflicting comments. In this paper we summarized in one table all the important information on drowning first aid to remind lifeguards, while on duty, what to do after the rescue of the victim. This was based on large rescue reports (1,2) and contributes to guide and develop standardized treatment protocols and also in comparing outcomes with different levels of severity of injury, thereby allowing lifeguards and medical staff to speak the same language - one team, one goal.

Four learning objectives

1. How to recognize drowning severity?
2. How to treat accordingly to severity?
3. When lifeguard should seek for medical advice or support?
4. To reassure lifeguards confidence in front of the public.

References:

1. Szpilman D. near-drowning and drowning classification: a proposal to stratify mortality based on the analysis of 1,831 cases. chest 1997; 112:660-5.
2. Szpilman D, Elmann J & Cruz-Filho FES; Drowning Classification: A Revalidation study based on the analysis of 930 cases over 10 Years; World Congress on Drowning, Netherlands 2002, Book of Abstracts, ISBN:90-6788-280-01, pg 66.
3. www.ilsf.org – ILS Policy Number 5, Medical Statement, Who needs further medical help after rescue from the water. Principal author: Ian Mackie; Approved by the ILS Board of Directors: December 1, 2000.

Principal Author: Dr David Szpilman, MD

Medical Staff of Drowning Resuscitation Center and Aeromedical Helicopter - Fire Department of Rio de Janeiro (CBMERJ) – GMAR; Head of Adult Intensive Care Unit - Hospital Municipal Miguel Couto; Member of Brazilian National Resuscitation Council; Founder, Former President and Medical Director of Brazilian Life Saving Society – SOBRASA; BOD Member and Medical Committee of International Life-Saving Federation; BOD Member of Sports Medical Council;

Av. das Américas 3555, Bloco 2, sala 302. Barra da Tijuca - RJ - Brazil 22631-004

Phones 055 21 99983951

Fone/Fax 055 21 33262378 or 24307168

Email: <david@szpilman.com>

Internet: <www.szpilman.com>

Grade	Frequency (%)	Signs	Treatment on land (basic and advanced life support)	Mortality on site (%)	Need of hospitalization	Mortality in hospital (%)
Rescue	0 to 6.4	Conscious, normal auscultation, no coughing, foam or difficulty in breathing	Evaluation and release from the accident site without further medical care if no other disease associated.	0	0%	0
1	56 to 65	Normal pulmonary auscultation, coughing, no foam in mouth or nose	Rest, warm and calm the victim. No further medical care.	0	2.9%	0
2	19 to 20	Rales in some pulmonary fields, small amount of foam in mouth or nose	5 liter oxygen per minute by nasal cannula; warm and calm the victim; recovery position, hospitalization	0,6 to 1,2	14.8%	4,0
3	3 to 3.3	Acute pulmonary edema with large amount of foam in mouth or nose, normotension, palpable radial pulse	15 liters oxygen per minute by non-rebreathing mask, intubation when possible; recovery position if unconscious; hospitalization.	3.6 to 5,2	44.8%	11,5
4	1.1 to 2	Acute pulmonary edema with large amount of foam in mouth or nose, hypotension verify by no radial pulse but palpable carotid pulse	Grade 3 treatment extended by crystalloid infusion via peripheral vein until restoration of Systolic Arterial Pressure > 90mmHg; hospitalization	19,4 to 22	88.9%	19,4
5	1 to 1.5	Isolated respiratory arrest	Mouth-to-mouth (when possible with 15 liters of oxygen) until restoration of normal breathing, then treat as grade 4.	31 to 44	100%	33,3
6	10 to 12	Cardiopulmonary arrest	Start CPR; insert tracheal tube, obtain venous access, give epinephrine 0,1 mg each 3 minute; ecg monitor and defibrillate if necessary. After successful CPR: follow grade 4.	88 to 93	100%	43,5
<i>Dead body</i>	----	Confirm submersion time over 1 hour, or obvious rigor mortis, putrefaction or dependent lividity	Do not start BLS or ALS; arrange morgue.	100	---	100

Should always receive medical attention (ambulance and/or hospitalization): Grades 2 to 6 and any patient in whom a serious condition is suspected such as heart attack, spinal injury, trauma, brief lost of consciousness, asthma, epilepsy, stinger, intoxication, delirium etc.

Should check before release to home: No cough; Normal breathing; Normal pulse (strength and rate); Normal color and skin perfusion; No shivering; and Fully conscious, awake and alert.

There is always a risk, although rare, of delayed lung complications: All victims should therefore be warned that if they later develop cough, breathlessness, fever or any other worrying symptom, they should seek medical advice immediately.

Table 1. Evaluation of 2,678 drowning severity and treatment decision scheme at the rescue site (Based on the assistance of 87,339 rescues) (1,2) and the indications needs of further medical help after rescue (3).